



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



THE GIFT OF
PROF. ALEXANDER ZIWET

QA
372
.081
1894

849

Alexander Zivox 3.7
EXAMPLES

OF

DIFFERENTIAL EQUATIONS

WITH

RULES FOR THEIR SOLUTION.

BY

GEORGE A. OSBORNE, S.B.

PROFESSOR OF MATHEMATICS IN THE MASSACHUSETTS INSTITUTE
OF TECHNOLOGY.



BOSTON, U.S.A. :
PUBLISHED BY GINN & COMPANY.

1894.



Entered, according to Act of Congress, in the year 1886, by
GEORGE A. OSBORNE,
in the Office of the Librarian of Congress, at Washington.

J. S. CUSHING & Co., PRINTERS, BOSTON.

From the
Estate of
T. C. Smith
3-18-30

PREFACE.

THIS work has been prepared to meet a want felt by the author in a practical course on the subject, arranged for advanced students in Physics. It is intended to be used in connection with lectures on the theory of Differential Equations and the derivation of the methods of solution.

Many of the examples have been collected from standard treatises, but a considerable number have been prepared by the author to illustrate special difficulties, or to provide exercises corresponding more nearly with the abilities of average students. With few exceptions they have all been tested by use in the class-room.

G. A. OSBORNE.

Boston, Feb 1, 1886.

Q 6-30-30 H.C.M.

CONTENTS.

CHAPTER I.

DEFINITIONS. — DERIVATION OF THE DIFFERENTIAL EQUATION FROM THE COMPLETE PRIMITIVE.

	PAGE.
Definitions	1
Derivation of differential equations of the first order	1
Derivation of differential equations of higher orders	2

SOLUTION OF DIFFERENTIAL EQUATIONS.

CHAPTER II.

DIFFERENTIAL EQUATIONS OF FIRST ORDER AND DEGREE BETWEEN TWO VARIABLES.

Form, $XYdx + X'Y'dy = 0$	4
Homogeneous equations	5
Form, $(ax + by + c)dx + (a'x + b'y + c')dy = 0$	5
Linear form, $\frac{dy}{dx} + Py = Q$	6
Form, $\frac{dy}{dx} + Py = Qy^n$	7

CHAPTER III.

EXACT DIFFERENTIAL EQUATIONS. — INTÉGRATING FACTORS.

Solution of exact differential equations	8
--	---

	PAGE.
Solution by means of an integrating factor in the following cases :	
Homogeneous equations	9
Form, $f_1(xy)y dx + f_2(xy)x dy = 0$	10
When $\frac{dM}{dy} \frac{dN}{dx} = \phi(x)$, or $\frac{dN}{dx} \frac{dM}{dy} = \psi(y)$	10

CHAPTER IV.

DIFFERENTIAL EQUATIONS OF FIRST ORDER AND DEGREE WITH THREE
VARIABLES.

Condition for a single primitive, and method of solution	12
--	----

CHAPTER V.

DIFFERENTIAL EQUATIONS OF THE FIRST ORDER OF HIGHER DEGREES.

Equations which can be solved with respect to p	14
Equations which can be solved with respect to y	14
Equations which can be solved with respect to x	15
Homogeneous equations	16
Clairaut's form, $y = px + f(p)$	16

CHAPTER VI.

SINGULAR SOLUTIONS.

Method of deriving the singular solution either from the complete primitive or from the differential equation.....	17
---	----

DIFFERENTIAL EQUATIONS OF HIGHER ORDERS.

CHAPTER VII.

LINEAR DIFFERENTIAL EQUATIONS.

Linear equations with constant coefficients and second member zero..	19
Linear equations with constant coefficients and second member not zero,	22
A special form of linear equations with variable coefficients	24

CONTENTS.

vii

CHAPTER VIII.

SPECIAL FORMS OF DIFFERENTIAL EQUATIONS OF HIGHER ORDERS.

	PAGE.
Form, $\frac{d^n y}{dx^n} = X$	25
Form, $\frac{d^2 y}{dx^2} = Y$	25
Equations not containing y directly	26
Equations not containing x directly	27

CHAPTER IX.

SIMULTANEOUS DIFFERENTIAL EQUATIONS.

Simultaneous equations of first order	28
Simultaneous equations of higher orders	30

CHAPTER X.

Geometrical applications	32
<hr/>	
ANSWERS TO EXAMPLES	35

EXAMPLES OF DIFFERENTIAL EQUATIONS.



CHAPTER I.

DEFINITIONS. DERIVATION OF THE DIFFERENTIAL EQUATION FROM THE COMPLETE PRIMITIVE.

1. A *differential equation* is an equation containing differentials or differential coefficients.

The *solution* of a differential equation is the determination of another equation free from differentials or differential coefficients, from which the former may be derived by differentiation.

The *order* of a differential equation is that of the highest differential coefficient it contains; and its *degree* is that of the highest power to which this highest differential coefficient is raised, after the equation is freed from fractions and radicals.

The solution of a differential equation requires one or more integrations, each of which introduces an arbitrary constant. The most general solution of a differential equation of the n th order contains n arbitrary constants, whatever may be its degree. This general solution is called the *complete primitive* of the given differential equation.

2. To derive a differential equation of the first order from its complete primitive.

Differentiate the primitive; and if the arbitrary constant has disappeared, the result is the required differential equation. If not, the elimination of this constant between the two equations will give the differential equation.

2 DERIVATION OF THE DIFFERENTIAL EQUATION.

3. Form the differential equations of the first order of which the following are the complete primitives, c being the arbitrary constant :

1. $\log(xy) + x = y + c.$
2. $(1 + x^2)(1 + y^2) = cx^2.$
3. $\cos y = c \cos x.$
4. $y = ce^{-\tan^{-1}x} + \tan^{-1}x - 1.$
5. $y = (cx + \log x + 1)^{-1}.$
6. $y = cx + c - c^3.$
7. $(y + c)^2 = 4ax.$
8. $y^2 \sin^2 x + 2cy + c^2 = 0.$
9. $e^{2y} + 2cxe^y + c^2 = 0.$

4. To derive a differential equation of the second order from its complete primitive.

Differentiate the primitive twice successively, and eliminate, if necessary, the two arbitrary constants between the three equations.

5. Form the differential equations of the second order of which the following are the complete primitives, c_1 and c_2 being the arbitrary constants :

1. $y = c_1 \cos(ax + c_2).$
2. $y = c_1 e^{ax} + c_2 e^{-ax}.$
3. $y = (c_1 + c_2 x)e^{ax}.$
4. $y = c_1 x^3 + \frac{c_2}{x}.$
5. $y = c_1 \sin nx + c_2 \cos nx + \frac{\cos ax}{n^2 - a^2}.$

6. The preceding process may be extended to the derivation of equations of higher orders from their primitives.

7. Form the differential equations of the third order of which the following are the complete primitives :

1. $y = c_1 e^{2x} + c_2 e^{3x} + c_3 e^x.$

2. $ye^x = c_1 e^{2x} + c_2 \sin x \sqrt{2} + c_3 \cos x \sqrt{2}.$

3. $y = \left(c_1 + c_2 x + \frac{x^2}{2} \right) e^x + c_3.$

Form the differential equations of the fourth order of which the following are the complete primitives :

4. $y = (c_1 + c_2 x + c_3 x^2) e^x + c_4.$

5. $x^3 + a^4 y = c_1 e^{ax} + c_2 e^{-ax} + c_3 \sin ax + c_4 \cos ax.$

CHAPTER II.

DIFFERENTIAL EQUATIONS OF THE FIRST ORDER AND FIRST DEGREE BETWEEN TWO VARIABLES.

General Form, $Mdx + Ndy = 0$,

where M, N , are each functions of x and y .

8. Form, $XYdx + X'Y'dy = 0$,

where X, X' , are functions of x alone, and Y, Y' , functions of y alone.

Divide so as to separate the variables, and integrate each part separately.

9. Solve the following equations :

1. $(1+x)ydx + (1-y)xdy = 0$.

2. $(x^2 - yx^2) \frac{dy}{dx} + y^2 + xy^2 = 0$.

3. $\frac{dy}{dx} = \frac{1+y^2}{(1+x^2)xy}$.

4. $a\left(x\frac{dy}{dx} + 2y\right) = xy\frac{dy}{dx}$.

5. $(1+y^2)dx = (y + \sqrt{1+y^2})(1+x^2)^{\frac{1}{2}}dy$.

6. $\sin x \cos y dx = \cos x \sin y dy$.

7. $\sec^2 x \tan y dx + \sec^2 y \tan x dy = 0$.

8. $\sec^2 x \tan y dy + \sec^2 y \tan x dx = 0$.

9. $\frac{dy}{dx} + \frac{1+y+y^2}{1+x+x^2} = 0$.

10. *Homogeneous equations.*

Substitute $y = vx$; in the resulting equation between v and x , the variables can be separated. (See Art. 8.)

11. Solve the following equations :

$$1. \quad (y - x)dy + ydx = 0.$$

$$2. \quad (2\sqrt{xy} - x)dy + ydx = 0.$$

$$3. \quad y^2 + x^2 \frac{dy}{dx} = xy \frac{dy}{dx}.$$

$$\times 4. \quad x \frac{dy}{dx} = y + \sqrt{x^2 + y^2}.$$

$$5. \quad x \cos \frac{y}{x} \cdot \frac{dy}{dx} = y \cos \frac{y}{x} - x.$$

$$\times 6. \quad (8y + 10x)dx + (5y + 7x)dy = 0.$$

$$\times 7. \quad (x + y) \frac{dy}{dx} = y - x.$$

$$8. \quad x \cos \frac{y}{x} (ydx + xdy) = y \sin \frac{y}{x} (xdy - ydx).$$

$$9. \quad x + y \frac{dy}{dx} = my.$$

$$(1), m < 2; \quad (2), m = 2; \quad (3), m > 2.$$

$$10. \quad [(x^2 - y^2) \sin \alpha + 2xy \cos \alpha - y \sqrt{x^2 + y^2}] \frac{dy}{dx} \\ = 2xy \sin \alpha - (x^2 - y^2) \cos \alpha + x \sqrt{x^2 + y^2}.$$

12. Form,

$$(ax + by + c)dx + (a'x + b'y + c')dy = 0.$$

$$\text{Substitute} \quad x = x' + \alpha, \quad y = y' + \beta,$$

and determine the constants α, β , so that the new equation between x' and y' may be homogeneous. (See Art. 10.)

This method fails when $\frac{a}{a'} = \frac{b}{b'}$. In this case put $ax + by = z$, and obtain a new equation between x and z or between y and z ; the variables can then be separated.

13. Solve the following equations :

$$1. \quad (3y - 7x + 7)dx + (7y - 3x + 3)dy = 0.$$

$$2. \quad (4x + 2y - 1)\frac{dy}{dx} + 2x + y + 1 = 0.$$

$$3. \quad \frac{dy}{dx} = \frac{7y + x + 2}{3x + 5y + 6}.$$

$$4. \quad (2y + x + 1)dx = (2x + 4y + 3)dy.$$

$$5. \quad 2x - y + 1 + (x + y - 2)\frac{dy}{dx} = 0.$$

14. *Linear Form,* $\frac{dy}{dx} + Py = Q,$

where P, Q , are independent of y .

Solution,
$$y = e^{-\int P dx} \left(\int Q e^{\int P dx} dx + c \right).$$

15. Solve the following equations :

$$1. \quad x \frac{dy}{dx} - ay = x + 1.$$

$$2. \quad x(1 - x^2)dy + (2x^2 - 1)ydx = ax^2dx.$$

$$3. \quad (1 - x^2)^2 \frac{dy}{dx} + y\sqrt{1 - x^2} = x + \sqrt{1 - x^2}.$$

$$4. \quad \frac{dy}{dx} + y \cos x = \frac{1}{2} \sin 2x.$$

$$5. \quad (1 + y^2)dx = (\tan^{-1}y - x)dy.$$

$$6. \quad \sqrt{a^2 + x^2} \left(1 - \frac{dy}{dx} \right) = x + y.$$

$$7. \quad (1 + x^2) dy + \left(xy - \frac{1}{x} \right) dx = 0.$$

$$8. \quad \frac{dy}{dx} + y \frac{d\phi}{dx} = \phi \frac{d\phi}{dx},$$

where ϕ is a function of x alone.

$$16. \text{ Form, } \frac{dy}{dx} + Py = Qy^n,$$

where P, Q , are independent of y .

Divide by y^n , and substitute $z = y^{-n+1}$. The new equation between z and x will be linear. (See Art. 14.)

17. Solve the following equations :

$$1. \quad (1 - x^2) \frac{dy}{dx} - xy = axy^2.$$

$$2. \quad 3y^2 \frac{dy}{dx} - ay^3 = x + 1.$$

$$3. \quad \frac{dy}{dx} = 2xy (ax^2y^2 - 1).$$

$$4. \quad \frac{dy}{dx} (x^2y^3 + xy) = 1.$$

$$5. \quad \frac{dy}{dx} + y \cos x = y^n \sin 2x.$$

$$6. \quad (y \log x - 1) y dx = x dy.$$

$$7. \quad ax^2y^n dy + y dx = 2x dy.$$

$$8. \quad y - \cos x \frac{dy}{dx} = y^2 \cos x (1 - \sin x).$$

$$9. \quad y \frac{dy}{dx} + by^2 = a \cos x.$$

CHAPTER III.

EXACT DIFFERENTIAL EQUATIONS AND INTEGRATING FACTORS.

18. $Mdx + Ndy$ is an exact differential when

$$\frac{dM}{dy} = \frac{dN}{dx}. \quad (1)$$

The solution of

$$Mdx + Ndy = 0, \quad \text{in this case is}$$

$$\int Mdx + \int \left(N - \frac{d}{dy} \int Mdx \right) dy = c,$$

or
$$\int Ndy + \int \left(M - \frac{d}{dx} \int Ndy \right) dx = c.$$

In integrating with respect to x , y is regarded as constant, and in integrating with respect to y , x is regarded as constant.

19. Solve the following equations after applying the condition (1) for an exact differential :

1. $(x^3 + 3xy^2) dx + (y^3 + 3x^2y) dy = 0.$

2. $(x^2 - 4xy - 2y^2) dx + (y^2 - 4xy - 2x^2) dy = 0.$

3. $\left(1 + \frac{y^2}{x^2} \right) dx - \frac{2y}{x} dy = 0.$

4. $\frac{2x dx}{y^3} + \left(\frac{1}{y^2} - \frac{3x^2}{y^4} \right) dy = 0.$

5. $x dx + y dy + \frac{x dy - y dx}{x^2 + y^2} = 0.$
6. $\frac{dx}{\sqrt{x^2 + y^2}} + \left(1 - \frac{x}{\sqrt{x^2 + y^2}}\right) \frac{dy}{y} = 0.$
7. $\left(x + \frac{1}{\sqrt{y^2 - x^2}}\right) dx + \left(y - \frac{x}{y \sqrt{y^2 - x^2}}\right) dy = 0.$
8. $(1 + e^{\frac{x}{y}}) dx + e^{\frac{x}{y}} \left(1 - \frac{x}{y}\right) dy = 0.$
9. $e^{\sqrt{x^2 + y^2 + 2x}} dx + 2ye^{\sqrt{x^2 + y^2 + 2x}} dy = 0.$
10. $(m dx + n dy) \sin(mx + ny) = (n dx + m dy) \cos(nx + my).$
11. $\frac{x dx + y dy}{\sqrt{1 + x^2 + y^2}} + \frac{y dx - x dy}{x^2 + y^2} = 0.$
12. $\frac{x^a dy - ayx^{a-1} dx}{by^2 - gx^{2a}} + x^{a-1} dx = 0.$
 (1), $g > 0$; (2), $g < 0$ and $= -k$; (3), $g = 0$;
 (4), $a = 0$; (5), $b = 0.$

20. When $Mdx + Ndy$ is not an exact differential, it may sometimes be made exact by multiplying by a factor, called an *integrating factor*. The following are some of the cases where this is possible.

21. When $Mdx + Ndy$ is homogeneous, $\frac{1}{Mx + Ny}$ is an integrating factor. This fails when $Mx + Ny = 0$, but in that case the solution is $y = cx$.

22. Solve the following equations by means of an integrating factor:

1. $(x^2 + 2xy - y^2) dx = (x^2 - 2xy - y^2) dy.$
2. $\frac{dx}{x} + \frac{dy}{y} + 2\left(\frac{dx}{y} - \frac{dy}{x}\right) = 0.$

$$3. \quad (x^2 y^2 + xy^3) dx - (x^3 y + x^2 y^2) dy = 0.$$

$$4. \quad x^3 dx + (3x^2 y + 2y^3) dy = 0.$$

$$5. \quad (x\sqrt{x^2 + y^2} - x^2) dy + (xy - y\sqrt{x^2 + y^2}) dx = 0$$

(See Art. 11 for other examples.)

$$23. \text{ Form, } f_1(xy)y dx + f_2(xy)x dy = 0.$$

$\frac{1}{Mx - Ny}$ is an integrating factor. This fails when

$$Mx - Ny = 0,$$

but in that case the solution is $xy = c$.

Another method of solving is to put $xy = v$, and obtain an equation between x and v or between y and v . The variables can then be separated.

24. Solve the following equations by means of an integrating factor :

$$1. \quad (1 + xy)y dx + (1 - xy)x dy = 0.$$

$$2. \quad (x^2 y^2 + xy)y dx + (x^2 y^2 - 1)x dy = 0.$$

$$3. \quad (x^3 y^3 + 1)(x dy + y dx) + (x^2 y^2 + xy)(y dx - x dy) = 0.$$

$$4. \quad (\sqrt{xy} - 1)x dy - (\sqrt{xy} + 1)y dx = 0.$$

$$5. \quad (y + y\sqrt{xy})dx + (x + x\sqrt{xy})dy = 0.$$

$$6. \quad e^{xy}(x^2 y^2 + xy)(x dy + y dx) + y dx - x dy = 0.$$

$$7. \quad xy[1 + \cot(xy)](x dy + y dx) + x dy - y dx = 0.$$

$$25. \text{ When } \frac{\frac{dM}{dy} - \frac{dN}{dx}}{N} = \phi(x),$$

then $e^{\int \phi(x) dx}$ is an integrating factor.

Or, when
$$\frac{\frac{dN}{dx} - \frac{dM}{dy}}{M} = \psi(y),$$

then $e^{\int \psi(y) dy}$ is an integrating factor.

26. Solve the following equations by means of an integrating factor :

1. $(x^2 + y^2 + 2x)dx + 2ydy = 0.$

2. $(3x^2 - y^2) \frac{dy}{dx} = 2xy.$

3. $\frac{dy}{dx} = \frac{x^2 + y^2}{2xy}.$

4. $[(1 - y)\sqrt{1 - x^2} - xy]dx + [1 - x^2 - x\sqrt{1 - x^2}]dy = 0.$

5. $(\cos x + 2y \sec y \sec^2 2x)dx + (\tan 2x \sec y - \sin x \tan y)dy = 0.$

6. $\sin(3x - 2y)(2dx - dy) + \sin(x - 2y)dy = 0.$

7. The Linear Equation

$$\frac{dy}{dx} + Py = Q,$$

where P and Q are independent of y .

CHAPTER IV.

DIFFERENTIAL EQUATIONS OF THE FIRST ORDER AND DEGREE CONTAINING THREE VARIABLES.

General form, $Pdx + Qdy + Rdz = 0$,

where P, Q, R , are each functions of x, y, z .

27. If the variables can be separated, solve by integrating the parts separately.

The equation is derivable from a single primitive only when the following condition is satisfied :

$$P\left(\frac{dQ}{dz} - \frac{dR}{dy}\right) + Q\left(\frac{dR}{dx} - \frac{dP}{dz}\right) + R\left(\frac{dP}{dy} - \frac{dQ}{dx}\right) = 0. \quad (1)$$

The solution may then be effected by first solving the equation with one of the parts Pdx, Qdy, Rdz , omitted, regarding x, y, z , respectively, constant.

Omitting Rdz , for example, we solve $Pdx + Qdy = 0$, regarding z constant, and introducing instead of an arbitrary constant of integration, Z , an undetermined function of z , which must be subsequently determined so that this primitive may satisfy the given differential equation. The equation of condition for determining Z will ultimately involve only Z and z .

28. Solve the following equations after applying the condition (1) for a single primitive :

1. $\frac{dx}{x-a} + \frac{dy}{y-b} + \frac{dz}{z-c} = 0.$

2. $(x - 3y - z)dx + (2y - 3x)dy + (z - x)dz = 0.$

3. $(y + z)dx + (z + x)dy + (x + y)dz = 0.$
4. $yzdx + zx dy + xy dz = 0.$
5. $(y + z)dx + dy + dz = 0.$
6. $ay^2z^2dx + bz^2x^2dy + cx^2y^2dz = 0.$
7. $zydx = zxdy + y^2dz.$
8. $(ydx + xdy)(a + z) = xydz.$
9. $(y + a)^2dx + zdy = (y + a)dz.$
10. $(y^2 + yz)dx + (xz + z^2)dy + (y^2 - xy)dz = 0.$
11. $(2x^2 + 2xy + 2xz^2 + 1)dx + dy + 2zdz = 0.$
12. $(x^2y - y^3 - y^2z)dx + (xy^2 - x^2z - x^3)dy + (xy^2 + x^2y)dz = 0.$

CHAPTER V.

DIFFERENTIAL EQUATIONS OF THE FIRST ORDER, OF A DEGREE ABOVE THE FIRST.

In what follows, p denotes $\frac{dy}{dx}$.

29. *When the equation can be solved with respect to p .*

The different values of p constitute so many differential equations of the first degree, which must be solved separately, using the same character for the arbitrary constant in all.

If the terms of each of these separate primitives be transposed to the first member, the product of these first members placed equal to zero will be the complete primitive.

30. Solve the following equations :

1. $p^2 - 5p + 6 = 0$.

2. $x^2 p^2 - a^2 = 0$.

✓ 3. $x p^2 - a = 0$.

4. $x p^2 = 1 - x$.

5. $x^2 p^2 + 3 x y p + 2 y^2 = 0$.

6. $p(p + y) = x(x + y)$.

7. $p^3 + 2 x p^2 - y^2 p^2 - 2 x y^2 p = 0$.

8. $p^3 - (x^2 + x y + y^2) p^2 + (x^3 y + x^2 y^2 + x y^3) p - x^3 y^3 = 0$.

9. $p^2 + 2 p y \cot x = y^2$.

31. *When the equation can be solved with respect to y .*

Differentiate, regarding p variable as well as x and y , and

substitute for dy , pdx . There will result a differential equation of the first degree between x and p . Solve this equation, and eliminate p between its primitive and the given equation.

32. Solve the following equations :

1. $x - yp = ap^2$.
2. $y = xp^2 + 2p$.
3. $(x + yp)^2 = a^2(1 + p^2)$.
- x 4. $y = xp + p - p^2$.
5. $(y - ap)^2 = 1 + p^2$.
6. $y = ap + bp^2$.
7. $x^2 + y = p^2$.
8. $y^2 = x^2(1 + p^2)$.
- x 9. $y = p^2 + 2p^3$.

33. *When the equation can be solved with respect to x .*

Differentiate, regarding p variable as well as x and y , and substitute for dx , $\frac{dy}{p}$. There will result a differential equation of the first degree between y and p . Solve this equation, and eliminate p between its primitive and the given equation.

34. Solve the following equations :

- x 1. $p^2y + 2px = y$.
2. $x = p + \log p$.
3. $p^2(x^2 + 2ax) = a^2$.
4. $x^2p^2 = 1 + p^2$.
5. $(x - ap)^2 = 1 + p^2$; also when $a = 1$.
6. $x = ap + bp^2$.
7. $my - nxp = yp^2$.

16 HOMOGENEOUS EQUATIONS. — CLAIRAUT'S FORM.

35. When the equation is homogeneous with respect to x and y .

Substitute $y = vx$. If the resulting equation between p and v can be solved with respect to v , the given equation comes under Art. 31 or Art. 33.

But if we can solve with respect to p , substitute for p , $v + x \frac{dv}{dx}$, and there will result a differential equation of the first degree between v and x .

36. Solve the following equations :

1. $xy^2(p^2 + 2) = 2py^3 + x^3.$
2. $(2p + 1)x^{\frac{1}{2}}y = x^{\frac{3}{2}}p^2 + 2y^{\frac{3}{2}}.$
3. $4x^2 = 3(3y - px)(y + px).$
4. $ds = \left(\frac{y}{2x}\right)^{\frac{1}{2}} dx + \left(\frac{x}{2y}\right)^{\frac{1}{2}} dy$, where $ds = \sqrt{1 + p^2} \cdot dx.$
5. $(nx + py)^2 = (1 + p^2)(y^2 + nx^2).$

37. Clairaut's Form,

$$y = px + f(p).$$

The solution is immediately obtained by substituting $p = c$.

38. Solve the following equations :

1. $y = px + \frac{m}{p}.$
2. $y = px + p - p^3.$
3. $y^2 - 2pxy - 1 = p^2(1 - x^2).$
4. $y = 2px + y^2p^3.$ Put $y^2 = y'.$
5. $ayp^3 + (2x - b)p = y.$ Put $y^2 = y'.$
6. $x^2(y - px) = yp^2.$ Put $y^2 = y', x^2 = x'.$
7. $e^{3x}(p - 1) + p^3e^{2y} = 0.$ Put $e^x = x', e^y = y'.$
8. $(px - y)(py + x) = h^2p.$ Put $y^2 = y', x^2 = x'.$

CHAPTER VI.

SINGULAR SOLUTIONS.

39. A *singular solution* of a differential equation is a solution which is not included in the complete primitive. Differential equations of the first degree have no singular solution. Those of higher degrees may have singular solutions, which may be derived either from the complete primitive, or directly from the differential equation.

40. Let $f(x, y, c) = 0$ be the complete primitive.

By differentiating, regarding c as the only variable, obtain $\frac{df}{dc} = 0$. If we eliminate c between this equation and the primitive, the result will be a singular solution, provided it satisfies the given differential equation.

41. Let $f(x, y, p) = 0$ be the given differential equation.

By differentiating, regarding p as the only variable, obtain $\frac{df}{dp} = 0$. If we eliminate p between this equation and the given differential equation, the result will be a singular solution, provided it satisfies the differential equation.

42. Derive the singular solution of the following equations, directly from the given equation, and also from the complete primitive :

1. $y = px + \frac{m}{p}.$

2. $y^2 - 2xyp + (1 + x^2)p^2 = 1.$

3. $p^3 - 4xyp + 8y^2 = 0$. Put $y = z^2$.
4. $y = (x - 1)p - p^2$.
5. $y(1 + p^2) = 2xp$.
6. $x^2p^2 - 2(xy - 2)p + y^2 = 0$.
7. $(y - xp)(mp - n) = mnp$.

DIFFERENTIAL EQUATIONS OF AN ORDER HIGHER THAN THE FIRST.



CHAPTER VII.

LINEAR DIFFERENTIAL EQUATIONS.

General Form,

$$\frac{d^n y}{dx^n} + X_1 \frac{d^{n-1} y}{dx^{n-1}} + X_2 \frac{d^{n-2} y}{dx^{n-2}} \dots + X_{n-1} \frac{dy}{dx} + X_n y = X,$$

the coefficients X_1, X_2, \dots, X_n and X being functions of x alone or constants.

43. Linear equations with *constant* coefficients and second member *zero* may be solved as follows :

Substitute in the given equation,

$$\frac{d^n y}{dx^n} = m^n, \quad \frac{d^{n-1} y}{dx^{n-1}} = m^{n-1}, \dots, \frac{dy}{dx} = m, \quad y = m^0 = 1.$$

There will result an equation of the n th degree in m , called the *auxiliary equation*. Find the n roots of this equation ; these roots will determine a series of terms expressing the complete value of y as follows, viz. For each real root m_1 , there will be a term $Ce^{m_1 x}$; for each pair of imaginary roots $a \pm b\sqrt{-1}$, a term $e^{ax}(A\sin bx + B\cos bx)$; each of the coefficients A, B, C , being an arbitrary constant if the corresponding root occur only once, but a polynomial $c_1 + c_2 x + c_3 x^2 \dots + c_r x^{r-1}$, if the root occur r times.

44. Roots of auxiliary equation, real and unequal.

Solve the following equations :

1. $\frac{d^2y}{dx^2} = a^2y.$

2. $\frac{d^2y}{dx^2} + 12y = 7\frac{dy}{dx}.$

3. $a\frac{d^2y}{dx^2} = \frac{dy}{dx}.$

4. $3\left(\frac{d^2y}{dx^2} + y\right) = 10\frac{dy}{dx}.$

5. $\frac{d^2y}{dx^2} + 4\frac{dy}{dx} = y.$

6. $ab\left(y + \frac{d^2y}{dx^2}\right) = (a^2 + b^2)\frac{dy}{dx}.$

7. $\frac{d^3y}{dx^3} = 4\frac{dy}{dx}.$

8. $\frac{d^3y}{dx^3} = \frac{d^2y}{dx^2} + 6\frac{dy}{dx}.$

9. $\frac{d^3y}{dx^3} = 7\frac{dy}{dx} - 6y.$

10. $\frac{d^4y}{dx^4} + 27y = 12\frac{d^2y}{dx^2}.$

11. $\frac{d^5y}{dx^5} - 2(a^2 + b^2)\frac{d^3y}{dx^3} + (a^2 - b^2)^2\frac{dy}{dx} = 0.$

45. Roots of auxiliary equation unequal, but not all real.

Solve the following equations :

1. $\frac{d^2y}{dx^2} + y = 0.$

2. $\frac{d^2y}{dx^2} - 6\frac{dy}{dx} + 13y = 0.$

$$3. \quad \frac{d^2 y}{dx^2} - 2a \frac{dy}{dx} + b^2 y = 0.$$

(1), when $a > b$; (2), when $a < b$.

$$4. \quad \frac{d^2 y}{dx^2} - 4ab \frac{dy}{dx} + (a^2 + b^2)^2 y = 0.$$

$$5. \quad \frac{d^2 y}{dx^2} - 2 \log a \frac{dy}{dx} + [1 + (\log a)^2] y = 0.$$

$$6. \quad \frac{d^3 y}{dx^3} + 2 \frac{dy}{dx} = 0.$$

$$7. \quad \frac{d^3 y}{dx^3} = y.$$

$$8. \quad \frac{d^3 y}{dx^3} + \frac{d^2 y}{dx^2} + \frac{dy}{dx} = 3y.$$

$$9. \quad \frac{d^4 y}{dx^4} = y.$$

$$10. \quad \frac{d^4 y}{dx^4} + 2 \frac{d^2 y}{dx^2} - 8y = 0.$$

$$11. \quad \frac{d^4 y}{dx^4} + 4a^4 y = 0.$$

$$12. \quad \frac{d^6 y}{dx^6} = y.$$

$$13. \quad \frac{d^6 y}{dx^6} = -y.$$

$$14. \quad \frac{d^8 y}{dx^8} = y.$$

46. *Auxiliary equation containing equal roots.*

Solve the following equations :

$$1. \quad \frac{d^2 y}{dx^2} - 2a \frac{dy}{dx} + a^2 y = 0.$$

$$2. \quad \frac{d^2 y}{dx^2} = 0.$$

$$3. \quad \frac{d^3 y}{dx^3} = 4 \frac{d^2 y}{dx^2}.$$

$$4. \quad \frac{d^3 y}{dx^3} - 3 \frac{d^2 y}{dx^2} + 4 y = 0.$$

$$5. \quad \frac{d^3 y}{dx^3} - \frac{d^2 y}{dx^2} - \frac{dy}{dx} + y = 0.$$

$$6. \quad \frac{d^4 y}{dx^4} + 2n^2 \frac{d^2 y}{dx^2} + n^4 y = 0.$$

$$7. \quad \frac{d^4 y}{dx^4} - 3 \frac{d^3 y}{dx^3} + 3 \frac{d^2 y}{dx^2} - \frac{dy}{dx} = 0.$$

$$8. \quad \frac{d^4 y}{dx^4} - 4 \frac{d^3 y}{dx^3} + 14 \frac{d^2 y}{dx^2} - 20 \frac{dy}{dx} + 25 y = 0, \quad \text{the first member of auxiliary equation being a perfect square.}$$

$$9. \quad \frac{d^n y}{dx^n} + \frac{d^{n-2} y}{dx^{n-2}} = 0.$$

47. Linear Equations with constant coefficients and second member *not* zero.

There are two methods of solution :

FIRST. METHOD OF VARIABLE PARAMETERS. — Solve the equation by Art. 43, regarding the second member as zero.

Supposing it to be of the n th order, this value of y will contain n arbitrary constants. Derive from it the successive differential coefficients, $\frac{dy}{dx}, \frac{d^2 y}{dx^2}, \dots, \frac{d^{n-1} y}{dx^{n-1}}$; then differentiate the values of $y, \frac{dy}{dx}, \frac{d^2 y}{dx^2}, \dots, \frac{d^{n-1} y}{dx^{n-1}}$, regarding the arbitrary constants alone as variable, and place these n results equal to zero, except the last, which put equal to the second member of the given equation. These n conditions will determine expressions for the n arbitrary constants, which are to be substituted in the original expression for y .

SECOND METHOD. — By successively differentiating the given equation, obtain, either directly or by elimination, a new differential equation of a higher order with the second member zero. Solve this by Art. 43, and determine the values of the superfluous constants so as to satisfy the given differential equation. In this last work of determining the superfluous constants all the other constants may be regarded as zero.

43. Solve the following equations :

$$1. \quad \frac{d^2 y}{dx^2} - 7 \frac{dy}{dx} + 12y = x.$$

$$2. \quad \frac{d^4 y}{dx^4} - 2 \frac{d^3 y}{dx^3} + 2 \frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} + y = a.$$

$$3. \quad \frac{d^2 y}{dx^2} - a^2 y = x + 1.$$

$$4. \quad \frac{d^3 y}{dx^3} - 2 \frac{d^2 y}{dx^2} + \frac{dy}{dx} = e^x.$$

$$5. \quad \frac{d^4 y}{dx^4} - a^4 y = x^3.$$

$$6. \quad \frac{d^2 y}{dx^2} + n^2 y = 1 + x + x^2.$$

$$7. \quad \frac{d^2 y}{dx^2} - 2a \frac{dy}{dx} + a^2 y = e^x; \quad \text{also when } a = 1.$$

$$8. \quad \frac{d^2 y}{dx^2} + n^2 y = \cos ax; \quad \text{also when } a = n.$$

$$9. \quad \frac{d^2 y}{dx^2} - 5 \frac{dy}{dx} + 6y = e^{nx}; \quad \text{also when } n = 2, \text{ or } n = 3.$$

$$10. \quad \frac{d^2 y}{dx^2} - 3 \frac{dy}{dx} + 2y = xe^{nx}.$$

$$11. \quad \frac{d^2 y}{dx^2} - 9 \frac{dy}{dx} + 20y = x^2 e^{3x}.$$

$$12. \quad \frac{d^2 y}{dx^2} + 4y = x \sin^2 x.$$

$$13. \quad \frac{d^4 y}{dx^4} + 2 \frac{d^2 y}{dx^2} + y = x^2 \cos ax; \quad \text{also when } a = 1.$$

$$14. \quad \frac{d^3 y}{dx^3} - 2 \frac{dy}{dx} + 4y = e^x \cos x.$$

49. Linear equations of the form

$$(a+bx)^n \frac{d^n y}{dx^n} + A_1(a+bx)^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots + A_{n-1}(a+bx) \frac{dy}{dx} + A_n y = X,$$

where A_1, A_2, \dots, A_n are constants, and X a function of x alone.

Put $a+bx=e^t$, and change the independent variable from x to t . The new differential equation between y and t will be linear with *constant* coefficients, and may be solved by Art. 47.

50. Solve the following equations :

$$1. \quad x^2 \frac{d^2 y}{dx^2} - x \frac{dy}{dx} = 3y.$$

$$2. \quad (x+a)^2 \frac{d^2 y}{dx^2} - 4(x+a) \frac{dy}{dx} + 6y = x.$$

$$3. \quad (a+bx)^2 \frac{d^2 y}{dx^2} + b(a+bx) \frac{dy}{dx} + b^2 y = 0.$$

$$4. \quad x^2 \frac{d^2 y}{dx^2} - x \frac{dy}{dx} + 2y = x \log x.$$

$$5. \quad (2x-1)^3 \frac{d^3 y}{dx^3} + (2x-1) \frac{dy}{dx} - 2y = 0.$$

$$6. \quad 16(x+1)^4 \frac{d^4 y}{dx^4} + 96(x+1)^3 \frac{d^3 y}{dx^3} + 104(x+1)^2 \frac{d^2 y}{dx^2} \\ + 8(x+1) \frac{dy}{dx} + y = x^2 + 4x + 3.$$

$$7. \quad x^4 \frac{d^4 y}{dx^4} + 6x^2 \frac{d^3 y}{dx^3} + 9x^2 \frac{d^2 y}{dx^2} + 3x \frac{dy}{dx} + y = (1 + \log x)^2.$$

$$8. \quad x^2 \frac{d^2 y}{dx^2} - (2m-1)x \frac{dy}{dx} + (m^2 + n^2)y = n^2 x^m \log x.$$

CHAPTER VIII.

SOME SPECIAL FORMS OF DIFFERENTIAL EQUATIONS OF HIGHER ORDERS.

51. Form, $\frac{d^n y}{dx^n} = X$, where X is a function of x alone.

The expression for y is found by integrating X successively n times with regard to x . Or solve by Art. 47.

52. Solve the following equations :

1. $x \frac{d^3 y}{dx^3} = 2.$

2. $\frac{d^4 y}{dx^4} = \frac{1}{(x+a)^2}.$

3. $\frac{d^n y}{dx^n} = x^m.$

4. $\frac{d^4 y}{dx^4} = x \cos x.$

5. $e^x \frac{d^4 y}{dx^4} + 4 \cos x = 0.$

6. $\frac{d^n y}{dx^n} = x e^x.$

7. $\frac{d^3 y}{dx^3} = \sin^3 x.$

53. Form, $\frac{d^2 y}{dx^2} = Y$, where Y is a function of y only.

Multiplying both members by $2 \frac{dy}{dx}$, and integrating, we have

$$\left(\frac{dy}{dx}\right)^2 = 2 \int Y dy + c_1. \quad \text{Therefore } x = \int \frac{dy}{(2 \int Y dy + c_1)^{\frac{1}{2}}} + c_2.$$

54. Solve the following equations :

$$1. \quad \frac{d^2 y}{dx^2} = a^2 y.$$

$$2. \quad \frac{d^2 y}{dx^2} = -a^2 y.$$

$$3. \quad y^3 \frac{d^2 y}{dx^2} = a.$$

$$4. \quad \frac{d^2 y}{dx^2} = e^{ny}.$$

$$5. \quad \frac{d^2 y}{dx^2} = \frac{1}{\sqrt{ay}}.$$

55. *Equations not containing y directly.*

By assuming the differential coefficient of the lowest order in the given equation equal to z , and consequently the other differential coefficients equal to the successive differential coefficients of z with respect to x , we shall obtain a new differential equation between z and x of a lower order than the given equation.

56. Solve the following equations :

$$1. \quad x \frac{d^2 y}{dx^2} + \frac{dy}{dx} = 0.$$

$$2. \quad \frac{d^2 y}{dx^2} = a^2 + b^2 \left(\frac{dy}{dx} \right)^2.$$

$$3. \quad \frac{dy}{dx} - x \frac{d^2 y}{dx^2} = f \left(\frac{d^2 y}{dx^2} \right).$$

$$4. \quad a^2 \left(\frac{d^2 y}{dx^2} \right)^2 = 1 + \left(\frac{dy}{dx} \right)^2.$$

$$5. \quad a^2 \left(\frac{d^2 y}{dx^2} \right)^2 = \left[1 + \left(\frac{dy}{dx} \right)^2 \right]^3.$$

$$6. \quad (1 + x^2) \frac{d^2 y}{dx^2} + 1 + \left(\frac{dy}{dx} \right)^2 = 0.$$

7. $(1 - x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} = 2.$
8. $2x \frac{d^3 y}{dx^3} \cdot \frac{d^2 y}{dx^2} = \left(\frac{d^2 y}{dx^2} \right)^2 - a^2.$
9. $\frac{d^3 y}{dx^3} \cdot \frac{d^2 y}{dx^2} = \left(1 - \frac{d^3 y}{dx^3} \right) \left[1 + \left(\frac{d^2 y}{dx^2} \right)^2 \right]^{\frac{1}{2}}.$
10. $\frac{d^3 y}{dx^3} \left(\frac{dy}{dx} \right)^3 = 1.$

57. *Equations not containing x directly.*

By assuming $\frac{dy}{dx} = z$, and consequently

$$\frac{d^2 y}{dx^2} = z \frac{dz}{dy}, \quad \frac{d^3 y}{dx^3} = z^2 \frac{d^2 z}{dy^2} + z \left(\frac{dz}{dy} \right)^2, \quad \text{etc.},$$

changing the independent variable from x to y , we shall obtain a new differential equation between z and y of a lower order than the given equation.

58. Solve the following equations :

1. $y \frac{d^2 y}{dx^2} + \left(\frac{dy}{dx} \right)^2 = 1.$
2. $y(1 - \log y) \frac{d^2 y}{dx^2} + (1 + \log y) \left(\frac{dy}{dx} \right)^2 = 0.$
3. $y \frac{d^2 y}{dx^2} + \left[\left(\frac{dy}{dx} \right)^2 + a^2 \left(\frac{d^2 y}{dx^2} \right)^2 \right]^{\frac{1}{2}} = \left(\frac{dy}{dx} \right)^2.$
4. $\left(\frac{dy}{dx} \right)^2 - y \frac{d^2 y}{dx^2} = \frac{dy}{dx} \cdot f \left[\left(\frac{dy}{dx} \right)^{-1} \cdot \frac{d^2 y}{dx^2} \right].$
5. $y \frac{d^2 y}{dx^2} - \left(\frac{dy}{dx} \right)^2 = y^2 \log y.$
6. $y^2 \left(\frac{d^2 y}{dx^2} \right)^2 + n \left(\frac{dy}{dx} \right)^4 = y^2 \frac{dy}{dx} \cdot \frac{d^3 y}{dx^3} + ny \left(\frac{dy}{dx} \right)^2 \frac{d^2 y}{dx^2}.$

CHAPTER IX.

SIMULTANEOUS DIFFERENTIAL EQUATIONS.

59. *Simultaneous differential equations of the first order.*

There should be n given equations between $n + 1$ variables. Selecting one of these for the independent variable, we may, by differentiating the given equations a sufficient number of times, eliminate all but one of the dependent variables and their differential coefficients. The resulting differential equation between two variables must be solved by the methods previously given, and from its primitive and the given equations may be obtained the values of the other dependent variables. The complete solution will consist of n equations containing n arbitrary constants.

In general, if we differentiate the given equations $n - 1$ times successively, we shall have in all n^2 equations, which are just sufficient for the elimination of $n - 1$ variables, together with their $n(n - 1)$ differential coefficients. Shorter processes for the elimination will frequently suggest themselves in special cases.

60. Solve the following simultaneous equations :

$$1. \quad \begin{cases} \frac{dx}{dt} + 4x + \frac{y}{4} = 0, \\ \frac{dy}{dt} + 3y - x = 0. \end{cases}$$

$$2. \quad -dx = \frac{dy}{3y + 4z} = \frac{dz}{2y + 5z}$$

$$3. \quad \frac{dx}{y - 7x} = \frac{-dy}{2x + 5y} = dt.$$

$$4. \quad \begin{cases} a \frac{dz}{dx} + n^2 y = e^x, \\ \frac{dy}{dx} + az = 0. \end{cases}$$

$$5. \quad \begin{cases} \frac{dx}{dt} + 5x + y = e^t, \\ \frac{dy}{dt} + 3y - x = e^{2t}. \end{cases}$$

$$6. \quad \frac{dx}{2y - 5x + e^t} = \frac{dy}{x - 6y + e^{2t}} = dt.$$

$$7. \quad \begin{cases} 4 \frac{dx}{dt} + 9 \frac{dy}{dt} + 44x + 49y = t, \\ 3 \frac{dx}{dt} + 7 \frac{dy}{dt} + 34x + 38y = e^t. \end{cases}$$

$$8. \quad \begin{cases} 4 \frac{dx}{dt} + 9 \frac{dy}{dt} + 11x + 31y = e^t, \\ 3 \frac{dx}{dt} + 7 \frac{dy}{dt} + 8x + 24y = e^{2t}. \end{cases}$$

$$9. \quad \begin{cases} 4 \frac{dx}{dt} + 9 \frac{dy}{dt} + 2x + 31y = e^t, \\ 3 \frac{dx}{dt} + 7 \frac{dy}{dt} + x + 24y = 3. \end{cases}$$

$$10. \quad \begin{cases} \frac{dx}{dt} + \frac{2}{t}(x - y) = 1, \\ \frac{dy}{dt} + \frac{1}{t}(x + 5y) = t. \end{cases}$$

$$11. \quad \begin{cases} t dx = (t - 2x) dt, \\ t dy = (tx + ty + 2x - t) dt. \end{cases}$$

$$12. \quad \begin{cases} \frac{dx}{dt} = ny - mz, \\ \frac{dy}{dt} = lz - nx, \\ \frac{dz}{dt} = mx - ly. \end{cases}$$

$$13. \quad \begin{cases} lt \frac{dx}{dt} = mn (y - z), \\ mt \frac{dy}{dt} = nl (z - x), \\ nt \frac{dz}{dt} = lm (x - y). \end{cases}$$

61. *Simultaneous differential equations of an order higher than the first.*

By differentiating the given equations a sufficient number of times, we may eliminate all but one of the dependent variables and their differential coefficients, and thus obtain a differential equation between two variables, which must be solved by the appropriate methods. Its primitive, together with the given equation, will enable us to determine the values of the other dependent variables. The general solution will contain a number of arbitrary constants equal to the sum of the highest orders of differential coefficients in the several given equations.

62. Solve the following simultaneous equations :

$$1. \quad \begin{cases} \frac{d^2 x}{dt^2} + n^2 x = 0, \\ \frac{d^2 y}{dt^2} - n^2 x = 0. \end{cases}$$

$$2. \quad \begin{cases} \frac{d^2 x}{dt^2} - 3x - 4y + 3 = 0, \\ \frac{d^2 y}{dt^2} + x + y + 5 = 0. \end{cases}$$

$$3. \quad \begin{cases} \frac{d^2 x}{dt^2} - 3x - 4y + 3 = 0, \\ \frac{d^2 y}{dt^2} + x - 8y + 5 = 0. \end{cases}$$

$$4. \quad \begin{cases} \frac{d^2 x}{dt^2} + n^2 y = 0, \\ \frac{d^2 y}{dt^2} - n^2 x = 0. \end{cases}$$

$$5. \quad \begin{cases} 2 \frac{d^2 y}{dx^2} - \frac{dz}{dx} - 4y = 2x, \\ 2 \frac{dy}{dx} + 4 \frac{dz}{dx} - 3z = 0. \end{cases}$$

$$6. \quad \begin{cases} \frac{d^3 y}{dx^3} + 4 \frac{d^2 z}{dx^2} + (5 - n^2) \frac{dy}{dx} + 2(1 - n^2)z = 0, \\ \frac{d^3 z}{dx^3} + 4 \frac{d^2 y}{dx^2} + (5 - n^2) \frac{dz}{dx} + 2(1 - n^2)y = 0. \end{cases}$$

$$7. \quad \begin{cases} \frac{d^4 y}{dx^4} - 4 \frac{d^3 z}{dx^3} + 4 \frac{d^2 y}{dx^2} - y = 0, \\ \frac{d^4 z}{dx^4} - 4 \frac{d^3 y}{dx^3} + 4 \frac{d^2 z}{dx^2} - z = 0. \end{cases}$$

CHAPTER X.

GEOMETRICAL EXAMPLES.

63. Expressions involved in the examples, p representing $\frac{dy}{dx}$, and q representing $\frac{d^2y}{dx^2}$.

$$\text{Subtangent} = \frac{y}{p}. \quad \text{Subnormal} = py.$$

$$\text{Normal} = y \sqrt{1 + p^2}. \quad \frac{ds}{dx} = \sqrt{1 + p^2}.$$

$$\text{Intercept of tangent on axis of } X = x - \frac{y}{p}.$$

$$\text{Intercept of tangent on axis of } Y = y - px.$$

$$\text{Radius of curvature} = \mp \frac{(1 + p^2)^{\frac{3}{2}}}{q}.$$

1. Find the curve whose subtangent varies as (is n times) the abscissa.
2. Find the curve whose subnormal is constant and equal to $2a$.
3. Find the curve whose normal is equal to the square of the ordinate.
4. Find the curve for which $s = mx^2$.
5. Find the curve for which $s^2 = y^2 - a^2$.

The *orthogonal trajectory* of a series of curves is a curve that intersects them all at right angles.

Describe the curves represented by the following equations, and find their orthogonal trajectories :

6. $y = mx$, m being the variable parameter.

7. $y^2 = 2ax - x^2$, a being the variable parameter.
8. $y^2 = 4ax$, a being the variable parameter.
9. $xy = k^2$, k being the variable parameter.
10. $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, b being the variable parameter.
11. $x^2 + m^2y^2 = m^2a^2$, a being the variable parameter.
12. $\frac{x^2}{b^2 + h^2} + \frac{y^2}{b^2} = 1$, b being the variable parameter.

The three following examples require the singular solution :

13. Find the curve such that the sum of the intercepts of the tangent on the axes of X and Y is constant and equal to a .
14. Find the curve such that the part of the tangent between the axes of X and Y is constant and equal to a .
15. Find the curve such that the area of the right triangle formed by the tangent with the axes of X and Y is constant and equal to a^2 .

The following examples require the solution of differential equations of the second order :

16. Find the curve such that the length of the arc measured from some fixed point of it is equal to the intercept of the tangent on the axis of X .
- x 17. Find the curve whose radius of curvature varies as (is n times) the cube of the normal.
- x 18. Find the curve whose radius of curvature is equal to the normal ; first, when the two have the same direction ; second, when they have opposite directions.
19. Find the curve whose radius of curvature is equal to twice the normal ; first, when the two have the same direction ; second, when they have opposite directions.

ANSWERS.



Art. 3. $\left(p = \frac{dy}{dx}\right)$

1. $y(1+x) + px(1-y) = 0.$
2. $(x^2 + 1)pxy = y^2 + 1.$
3. $\tan x = p \tan y.$
4. $(1+x^2)p + y = \tan^{-1}x.$
5. $(y \log x - 1)y = px.$
6. $y = px + p - p^3.$
7. $xp^2 = a.$
8. $p^2 + 2py \cot x = y^2.$
9. $x^2 p^2 = 1 + p^2.$

Art. 5.

1. $\frac{d^2 y}{dx^2} + a^2 y = 0.$
2. $\frac{d^2 y}{dx^2} - a^2 y = 0.$
3. $\frac{d^2 y}{dx^2} - 2a \frac{dy}{dx} + a^2 y = 0.$
4. $x^2 \frac{d^2 y}{dx^2} - x \frac{dy}{dx} = 3y.$
5. $\frac{d^2 y}{dx^2} + n^2 y = \cos ax.$

Art. 7.

1. $\frac{d^3 y}{dx^3} = 7 \frac{dy}{dx} - 6y.$
2. $\frac{d^3 y}{dx^3} + \frac{d^2 y}{dx^2} + \frac{dy}{dx} = 3y.$
3. $\frac{d^3 y}{dx^3} - 2 \frac{d^2 y}{dx^2} + \frac{dy}{dx} = e^x.$
4. $\frac{d^4 y}{dx^4} - 3 \frac{d^3 y}{dx^3} + 3 \frac{d^2 y}{dx^2} - \frac{dy}{dx} = 0.$
5. $\frac{d^4 y}{dx^4} - a^4 y = x^3.$

Art. 9.

1. $\log(xy) + x - y = c.$
2. $\frac{x+y}{xy} + \log \frac{y}{x} = c.$
3. $(1+x^2)(1+y^2) = cx^2.$
4. $x^2 y = ce^{\frac{y}{x}}.$

$$5. \log [(y + \sqrt{1+y^2}) \sqrt{1+y^2}] = \frac{x}{\sqrt{1+x^2}} + c.$$

$$6. \cos y = c \cos x.$$

$$8. \sin^2 x + \sin^2 y = c.$$

$$7. \tan x \tan y = c.$$

$$9. xy - 1 = c(x + y + 1).$$

Art. 11.

$$1. y = ce^{-\frac{x}{y}}.$$

$$5. x = ce^{-\sin \frac{y}{x}}.$$

$$2. y = ce^{-\sqrt{\frac{x}{y}}}.$$

$$6. (y+x)^2 (y+2x)^3 = c.$$

$$3. y = ce^{\frac{y}{x}}.$$

$$7. \log (x^2 + y^2) = 2 \tan^{-1} \frac{x}{y} + c. \quad r = c e^{-\varphi}$$

$$4. x^2 = c^2 + 2cy.$$

$$8. xy \cos \frac{y}{x} = c.$$

$$9. (1), \log (x^2 - mxy + y^2) + \frac{2m}{\sqrt{4-m^2}} \tan^{-1} \frac{2y-mx}{x\sqrt{4-m^2}} = c.$$

$$(2), x - y = ce^{\frac{x}{y-x}}.$$

$$(3), \frac{(2y - mx + x\sqrt{m^2 - 4})^{m - \sqrt{m^2 - 4}}}{(2y - mx - x\sqrt{m^2 - 4})^{m + \sqrt{m^2 - 4}}} = c.$$

$$10. y \sin a - x \cos a + \sqrt{x^2 + y^2} = c(x^2 + y^2).$$

Art. 13.

$$1. (y - x + 1)^2 (y + x - 1)^5 = c.$$

$$2. x + 2y + \log (2x + y - 1) = c.$$

$$3. x + 5y + 2 = c(x - y + 2)^4.$$

$$4. 4x - 8y = \log (4x + 8y + 5) + c.$$

$$5. \log [2(3x-1)^2 + (3y-5)^2] - \sqrt{2} \tan^{-1} \frac{\sqrt{2}(3x-1)}{3y-5} = c.$$

Art. 15.

1. $y = cx^a + \frac{x}{1-a} - \frac{1}{a}$.
2. $y = ax + cx\sqrt{1-x^2}$.
3. $y = \frac{x}{\sqrt{1-x^2}} + ce^{\frac{x}{\sqrt{1-x^2}}}$.
4. $y = \sin x - 1 + ce^{-\sin x}$.
5. $x = \tan^{-1}y - 1 + ce^{-\tan^{-1}y}$.
6. $(x + \sqrt{a^2 + x^2})y = a^2 \log(x + \sqrt{a^2 + x^2}) + c$.
7. $y\sqrt{1+x^2} = \log \frac{\sqrt{1+x^2} - 1}{x} + c$.
8. $y = ce^{-\phi} + \phi - 1$.

Art. 17.

1. $y = (c\sqrt{1-x^2} - a)^{-1}$.
2. $y^3 = ce^{ax} - \frac{x+1}{a} - \frac{1}{a^2}$.
3. $y = \left[ce^{2x^2} + \frac{a}{2}(2x^2 + 1) \right]^{-\frac{1}{2}}$.
4. $x = \frac{e^{\frac{y^2}{2}}}{(2-y^2)e^{\frac{y^2}{2}} + c}$.
5. $y^{-n+1} = ce^{(n-1)\sin x} + 2\sin x + \frac{2}{n-1}$.
6. $y = (cx + \log x + 1)^{-1}$.
7. $x = \frac{(n+2)y^2}{ay^{n+2} + c}$.
8. $y = \frac{\tan x + \sec x}{\sin x + c}$.
9. $(4b^2 + 1)y^2 = 2a(\sin x + 2b \cos x) + ce^{-2bx}$.

Art. 19.

1. $x^4 + 6x^2y^2 + y^4 = c$.
2. $x^3 - 6x^2y - 6xy^2 + y^3 = c$.
3. $x^2 - y^2 = cx$.
4. $x^2 - y^2 = cy^3$.
5. $x^2 + y^2 + 2\tan^{-1}\frac{y}{x} = c$.
6. $y^2 = c^2 - 2cx$.
7. $x^2 + y^2 + 2\sin^{-1}\frac{x}{y} = c$.
8. $x + ye^{\frac{x}{y}} = c$.

9. $e^x(x^2 + y^2) = c.$
10. $\cos(mx + ny) + \sin(nx + my) = c.$
11. $\sqrt{1 + x^2 + y^2} + \tan^{-1} \frac{x}{y} = c.$
12. (1), $\log \frac{x^a \sqrt{g} + y \sqrt{b}}{x^a \sqrt{g} - y \sqrt{b}} = \frac{2x^a \sqrt{bg}}{a} + c.$
- (2), $\tan^{-1} \frac{y \sqrt{b}}{x^a \sqrt{k}} + \frac{x^a \sqrt{bk}}{a} = c.$
- (3), $x^a \left(\frac{1}{a} - \frac{1}{by} \right) = c.$
- (4), $\frac{\sqrt{g} + y \sqrt{b}}{\sqrt{g} - y \sqrt{b}} = cx^{2\sqrt{bg}}.$
- (5), $\frac{x^a}{a} - \frac{y}{gx^a} = c.$

Art. 22.

1. $x^2 + y^2 = c(x + y).$
2. $x^2 - y^2 + xy = c.$
3. $y = cx.$
4. $x^2 + 2y^2 = c\sqrt{x^2 + y^2}.$
5. $y = cx.$

Art. 24.

1. $x = cye^{\frac{1}{xy}}.$
2. $y = ce^{xy}.$
3. $xy - \frac{1}{xy} = \log cy^2.$
4. $\frac{2}{\sqrt{xy}} = \log \frac{cx}{y}.$
5. $xy = c.$
6. $xye^{xy} = \log \frac{cy}{x}.$
7. $xy + \log \sin(xy) = \log \frac{cx}{y}.$

Art. 26.

1. $e^x(x^2 + y^2) = c.$
2. $x^2 - y^2 = cy^3.$
3. $x^2 - y^2 = cx.$
4. $y\sqrt{1 - x^2} + x(1 - y) = c.$

$$5. \sin x \cos y + y \tan 2x = c. \quad 6. \sin^2 x \sin 2(x - y) = c.$$

$$7. y = e^{-\int P dx} \left(\int Q e^{\int P dx} dx + c \right).$$

Art. 28.

$$1. (x - a)(y - b)(z - c) = c'. \quad 7. z = ce^{\frac{x}{y}}.$$

$$2. x^2 + 2y^2 - 6xy - 2xz + z^2 = c. \quad 8. xy = c(a + z).$$

$$3. yz + zx + xy = c. \quad 9. x = \frac{z}{y + a} + c.$$

$$4. xyz = c. \quad 10. y(x + z) = c(y + z).$$

$$5. e^x(y + z) = c. \quad 11. e^{x^2}(x + y + z^2) = c.$$

$$6. \frac{a}{x} + \frac{b}{y} + \frac{c}{z} = c'. \quad 12. \frac{y + z}{x} + \frac{z + x}{y} = c.$$

Art. 30.

$$1. (y - 2x + c)(y - 3x + c) = 0, \text{ or } (5x - 2y + c)^2 = x^2.$$

$$2. (y + c)^2 = a^2(\log x)^2. \quad 3. (y + c)^2 = 4ax.$$

$$4. (y + c)^2 = (\sqrt{x - x^2} + \sin^{-1} \sqrt{x})^2.$$

$$5. (xy + c)(x^2y + c) = 0.$$

$$6. (x^2 - 2y + c)[e^x(x + y - 1) + c] = 0.$$

$$7. (y + c)(y + x^2 + c)(xy + cy + 1) = 0.$$

$$8. (x^2 - 3y + c)(e^{\frac{x^2}{2}} + cy)(xy + cy + 1) = 0.$$

$$9. y^2 \sin^2 x + 2cy + c^2 = 0.$$

Art. 32.

$$1. \text{ Eliminate } p \text{ by means of } x = \frac{p}{\sqrt{1 - p^2}}(c + a \sin^{-1} p).$$

$$2. \text{ Eliminate } p \text{ by means of } x(p - 1)^2 = \log p^2 - 2p + c.$$

$$3. \text{ Eliminate } p \text{ by means of } x = \frac{p}{\sqrt{1 + p^2}} \left(c + \frac{a}{p} + a \tan^{-1} p \right).$$

$$4. y = cx + c - c^2.$$

$$5. x = a \log (ay \pm \sqrt{a^2 + y^2 - 1}) + \log (y \mp \sqrt{a^2 + y^2 - 1}) + c.$$

$$6. x \pm \sqrt{a^2 + 4by} = a \log (a \pm \sqrt{a^2 + 4by}) + c.$$

$$7. c(2y \pm x\sqrt{x^2 + y})^{\sqrt{17}} = \frac{\pm 4\sqrt{x^2 + y} - x(\sqrt{17} - 1)}{\pm 4\sqrt{x^2 + y} + x(\sqrt{17} + 1)}.$$

$$8. \left[y\sqrt{y^2 - x^2} - x^2 \log \frac{y + \sqrt{y^2 - x^2}}{x} \right]^2 = (y^2 + x^2 \log cx^2)^2.$$

$$9. 4(x+c)^3 + (x+c)^2 - 18y(x+c) - 27y^2 - 4y = 0.$$

Art. 34.

$$1. y^2 = 2cx + c^2.$$

$$2. x + 1 = \pm \sqrt{2y + c} + \log (\pm \sqrt{2y + c} - 1).$$

$$3. (e^{\frac{y}{a}} - ac)^2 = 2cxe^{\frac{y}{a}}.$$

$$4. e^{2y} + 2cxe^y + c^2 = 0.$$

$$5. 2y + c = \frac{ax^2 \pm x\sqrt{a^2 + x^2 - 1}}{a^2 - 1} + \log (x \pm \sqrt{a^2 + x^2 - 1});$$

$$\text{when } a=1, \quad 4y = x^2 - \log (cx^2).$$

$$6. b^2(6y + c)^2 + (6abx + a^3)(6y + c) - 3a^2x^2 - 16bx^3 = 0.$$

$$7. c(nx^2 + 2y^2 \pm x\sqrt{n^2x^2 + 4my^2})^n = [(2m - n)x \pm \sqrt{n^2x^2 + 4my^2}]^{2m}.$$

Art. 36.

$$1. (x^2 - y^2 + c)(x^2 - y^2 + cx^4) = 0.$$

$$2. (\sqrt{x} - \sqrt{y} + c)(\sqrt{x} - \sqrt{y} + cx) = 0.$$

$$3. 3x^4 + 6cxy + c^2 = 0.$$

$$4. (y - x)^{1 \pm \sqrt{2}} = c(\sqrt{y} + \sqrt{x})^2.$$

$$5. x^{2k} + 2cx^{k-1}y - c^2n = 0, \quad \text{where } k = \sqrt{\frac{n-1}{n}}.$$

Art. 38.

1. $y = cx + \frac{m}{c}.$

5. $4y^2 = 2c(2x - b) + ac^2.$

2. $y = cx + c - c^3.$

6. $y^2 = cx^2 + c^2.$

3. $(y - cx)^2 = 1 + c^2.$

7. $e^y = ce^x + c^3.$

4. $y^2 = cx + \frac{c^3}{8}.$

8. $y^2 - cx^2 = -\frac{ch^2}{c+1}.$

Art. 42.

Complete Primitives and Singular Solutions :

1. $y = cx + \frac{m}{c},$

$y^2 = 4mx.$

2. $(y - cx)^2 = 1 - c^2,$

$y^2 - x^2 = 1.$

3. $y = c(x - c)^2,$

$y = \frac{4x^3}{27}.$

4. $y = c(x - 1) - c^2,$

$4y = (x - 1)^2.$

5. $y^2 - 2cx + c^2 = 0,$

$y^2 = x^2.$

6. $(y - cx)^2 + 4c = 0,$

$xy = 1.$

7. $(y - cx)(mc - n) = mnc,$

$\left(\frac{x}{m}\right)^{\frac{1}{2}} \pm \left(\frac{y}{n}\right)^{\frac{1}{2}} = 1.$

Art. 44.

1. $y = c_1 e^{ax} + c_2 e^{-ax}.$

5. $ye^{2x} = c_1 e^{x\sqrt{5}} + c_2 e^{-x\sqrt{5}}.$

2. $y = c_1 e^{3x} + c_2 e^{4x}.$

6. $y = c_1 e^{\frac{ax}{b}} + c_2 e^{\frac{bx}{a}}.$

3. $y = c_1 e^{\frac{x}{a}} + c_2.$

7. $y = c_1 e^{2x} + c_2 e^{-2x} + c_3.$

4. $y = c_1 e^{3x} + c_2 e^{\frac{x}{3}}.$

8. $y = c_1 e^{3x} + c_2 e^{-2x} + c_3.$

9. $y = c_1 e^{2x} + c_2 e^{-3x} + c_3 e^x.$

10. $y = c_1 e^{3x} + c_2 e^{-3x} + c_3 e^{x\sqrt{3}} + c_4 e^{-x\sqrt{3}}.$

11. $y = c_1 e^{(a-b)x} + c_2 e^{(b-a)x} + c_3 e^{(a+b)x} + c_4 e^{-(a+b)x} + c_5.$

Art. 45.

1. $y = c_1 \sin x + c_2 \cos x$.
2. $y = e^{\beta x} (c_1 \sin 2x + c_2 \cos 2x)$.
3. When $a > b$, $y = e^{ax} (c_1 e^{x\sqrt{a^2-b^2}} + c_2 e^{-x\sqrt{a^2-b^2}})$;
when $a < b$, $y = e^{ax} (c_1 \sin x \sqrt{b^2-a^2} + c_2 \cos x \sqrt{b^2-a^2})$.
4. $y = e^{2abx} [c_1 \sin (a^2 - b^2)x + c_2 \cos (a^2 - b^2)x]$.
5. $y = a^x (c_1 \sin x + c_2 \cos x)$.
6. $y = c_1 \sin x \sqrt{2} + c_2 \cos x \sqrt{2} + c_3$.
7. $y = c_1 e^x + e^{-\frac{x}{2}} \left(c_2 \sin \frac{x\sqrt{3}}{2} + c_3 \cos \frac{x\sqrt{3}}{2} \right)$.
8. $ye^x = c_1 e^{2x} + c_2 \sin x \sqrt{2} + c_3 \cos x \sqrt{2}$.
9. $y = c_1 e^x + c_2 e^{-x} + c_3 \sin x + c_4 \cos x$.
10. $y = c_1 e^{x\sqrt{2}} + c_2 e^{-x\sqrt{2}} + c_3 \sin 2x + c_4 \cos 2x$.
11. $y = e^{ax} (c_1 \sin ax + c_2 \cos ax) + e^{-ax} (c_3 \sin ax + c_4 \cos ax)$.
12. $y = c_1 e^x + c_2 e^{-x} + (c_3 e^{\frac{x}{2}} + c_4 e^{-\frac{x}{2}}) \sin \frac{x\sqrt{3}}{2}$
 $+ (c_5 e^{\frac{x}{2}} + c_6 e^{-\frac{x}{2}}) \cos \frac{x\sqrt{3}}{2}$.
13. $y = c_1 \sin x + c_2 \cos x + e^{\frac{x\sqrt{3}}{2}} \left(c_3 \sin \frac{x}{2} + c_4 \cos \frac{x}{2} \right)$
 $+ e^{-\frac{x\sqrt{3}}{2}} \left(c_5 \sin \frac{x}{2} + c_6 \cos \frac{x}{2} \right)$.
14. $y = c_1 e^x + c_2 e^{-x} + c_3 \sin x + c_4 \cos x + e^{\frac{x}{\sqrt{2}}} \left(c_5 \sin \frac{x}{\sqrt{2}} + c_6 \cos \frac{x}{\sqrt{2}} \right)$
 $+ e^{-\frac{x}{\sqrt{2}}} \left(c_7 \sin \frac{x}{\sqrt{2}} + c_8 \cos \frac{x}{\sqrt{2}} \right)$.

Art. 46.

1. $y = (c_1 + c_2 x) e^{ax}$.
2. $y = c_1 + c_2 x$.
3. $y = c_1 e^{4x} + c_2 + c_3 x$.
4. $y = c_1 e^{-x} + (c_2 + c_3 x) e^{2x}$.

5. $y = c_1 e^{-x} + (c_2 + c_3 x) e^x.$
6. $y = (c_1 + c_2 x) \cos nx + (c_3 + c_4 x) \sin nx.$
7. $y = (c_1 + c_2 x + c_3 x^2) e^x + c_4.$
8. $y = e^x [(c_1 + c_2 x) \sin 2x + (c_3 + c_4 x) \cos 2x].$
9. $y = c_1 + c_2 x + c_3 x^2 \dots + c_{n-2} x^{n-3} + c_{n-1} \sin x + c_n \cos x.$

Art. 48.

1. $y = c_1 e^{3x} + c_2 e^{4x} + \frac{12x+7}{144}.$
2. $y = c_1 \sin x + c_2 \cos x + (c_3 + c_4 x) e^x + a.$
3. $y = c_1 e^{ax} + c_2 e^{-ax} - \frac{x+1}{a^2}.$
4. $y = \left(c_1 + c_2 x + \frac{x^2}{2} \right) e^x + c_3.$
5. $y = c_1 e^{ax} + c_2 e^{-ax} + c_3 \sin ax + c_4 \cos ax - \frac{x^3}{a^4}.$
6. $y = c_1 \sin nx + c_2 \cos nx + \frac{1+x+x^2}{n^2} - \frac{2}{n^4}.$
7. $y = (c_1 + c_2 x) e^{ax} + \frac{e^x}{(a-1)^2};$
when $a = 1, \quad y = \left(c_1 + c_2 x + \frac{x^2}{2} \right) e^x.$
8. $y = c_1 \sin nx + c_2 \cos nx + \frac{\cos ax}{n^2 - a^2};$
when $a = n, \quad y = c_1 \sin nx + c_2 \cos nx + \frac{x \sin nx}{2n}.$
9. $y = c_1 e^{2x} + c_2 e^{3x} + \frac{e^{nx}}{n^2 - 5n + 6};$
when $n = 2, \quad y = (c_1 - x) e^{2x} + c_2 e^{3x};$
when $n = 3, \quad y = c_1 e^{2x} + (c_2 + x) e^{3x}.$
10. $y = c_1 e^x + c_2 e^{2x} + \frac{x e^{nx}}{n^2 - 3n + 2} - \frac{(2n-3) e^{nx}}{(n^2 - 3n + 2)^2}.$

$$11. y = c_1 e^{4x} + c_2 e^{5x} + \frac{2x^2 + 6x + 7}{4} e^{3x}.$$

$$12. y = \left(c_1 - \frac{x^2}{16}\right) \sin 2x + \left(c_2 - \frac{x}{32}\right) \cos 2x + \frac{x}{8}.$$

$$13. y = (c_1 + c_2 x) \sin x + (c_3 + c_4 x) \cos x \\ + \left[\frac{x^2}{(a^2 - 1)^2} - \frac{4(5a^2 + 1)}{(a^2 - 1)^4} \right] \cos ax - \frac{8ax}{(a^2 - 1)^3} \sin ax;$$

$$\text{when } a = 1, \quad y = \left(c_1 + c_2 x + \frac{x^3}{12}\right) \sin x \\ + \left(c_3 + c_4 x - \frac{x^4}{48} + \frac{3x^2}{16}\right) \cos x.$$

$$14. y = c_1 e^{-2x} + \left(c_2 - \frac{x}{20}\right) e^x \cos x + \left(c_3 + \frac{3x}{20}\right) e^x \sin x.$$

Art. 50.

$$1. y = c_1 x^3 + \frac{c_2}{x}.$$

$$2. y = c_1(x + a)^2 + c_2(x + a)^3 + \frac{3x + 2a}{6}.$$

$$3. y = c_1 \sin \log(a + bx) + c_2 \cos \log(a + bx).$$

$$4. y = x(c_1 \sin \log x + c_2 \cos \log x + \log x).$$

$$5. y = (2x - 1) \left[c_1 + c_2 (2x - 1)^{\frac{\sqrt{3}}{2}} + c_3 (2x - 1)^{-\frac{\sqrt{3}}{2}} \right].$$

$$6. y = [c_1 + c_2 \log(x + 1)] \sqrt{x + 1} + \frac{c_3 + c_4 \log(x + 1)}{\sqrt{x + 1}} \\ + \frac{x^2 + 52x + 51}{225}.$$

$$7. y = (c_1 + c_2 \log x) \sin \log x + (c_3 + c_4 \log x) \cos \log x + (\log x)^2 \\ + 2 \log x - 3.$$

$$8. y = x^m (c_1 \sin \log x^n + c_2 \cos \log x^n + \log x).$$

Art. 52.

$$1. y = c_1 + c_2 x + c_3 x^2 + x^2 \log x.$$

$$2. y = c_1 + c_2 x + c_3 x^2 + c_4 x^3 - (x + a)^2 \log \sqrt{x + a}.$$

3. $y = c_1 + c_2x + c_3x^2 \dots + c_nx^{n-1} + \frac{mx^{m+n}}{m+n}.$
4. $y = c_1 + c_2x + c_3x^2 + c_4x^3 + x \cos x - 4 \sin x.$
5. $y = c_1 + c_2x + c_3x^2 + c_4x^3 + e^{-x} \cos x.$
6. $y = c_1 + c_2x + c_3x^2 \dots + c_nx^{n-1} + (x-n)e^x.$
7. $y = c_1 + c_2x + c_3x^2 + \frac{7 \cos x}{9} - \frac{\cos^3 x}{27}.$

Art. 54.

1. $ax = \log(y + \sqrt{y^2 + c_1}) + c_2,$ or $y = c'_1 e^{ax} + c'_2 e^{-ax}.$
2. $ax = \sin^{-1} \frac{y}{c_1} + c_2,$ or $y = c_1 \sin(ax + c_2).$
3. $(c_1x + c_2)^2 + a = c_1y^2.$
4. $x\sqrt{2n} = c_1 \log \frac{\sqrt{c_1^2 e^{ny} + 1} - 1}{\sqrt{c_1^2 e^{ny} + 1} + 1} + c_2.$
5. $3x = 2a^{\frac{1}{2}}(y^{\frac{1}{2}} - 2c_1)(y^{\frac{1}{2}} + c_1)^{\frac{1}{2}} + c_2.$

Art. 56.

1. $y = c_1 \log x + c_2.$
2. $b^2y = \log \sec[ab(x + c_1)] + c_2.$
3. $y = \frac{c_1x^2}{2} + xf(c_1) + c_2.$
4. $\frac{2y}{a} = c_1 e^{\frac{x}{a}} + c_1^{-1} e^{-\frac{x}{a}} + c_2.$
5. $(x + c_1)^2 + (y + c_2)^2 = a^2.$
6. $y = c_1x + (c_1^2 + 1) \log(x - c_1) + c_2.$
7. $y = c_1 \sin^{-1}x + (\sin^{-1}x)^2 + c_2.$
8. $y = \frac{4}{15c_1}(x + c_1^2a^2)^{\frac{5}{2}} + c_2x + c_3.$

$$9. \quad 12y = w^3 + c_1 w - 6w \log w + c_2, \quad \text{where } w = x + c_3.$$

$$10. \quad 2y\sqrt{c_1} = w\sqrt{w^2 + c_1^2} + c_1^2 \log(w + \sqrt{w^2 + c_1^2}) + c_2, \\ \text{where } w = x + c_3.$$

Art. 58.

$$1. \quad y^2 = x^2 + c_1 x + c_2.$$

$$4. \quad c_1 x = \log[c_1 y + f(c_1)] + c_2.$$

$$2. \quad \log y - 1 = \frac{1}{c_1 x + c_2}.$$

$$5. \quad \log y = c_1 e^x + c_2 e^{-x}.$$

$$3. \quad c_1 y = c_2 e^{c_1 x} - \sqrt{1 + a^2 c_1^2}.$$

$$6. \quad y^{n+1} = c_1 e^{ax} + c_2.$$

Art. 60.

$$1. \quad \begin{cases} 2x = (2c_2 - c_1 - c_2 t) e^{-\frac{7t}{2}}, \\ y = (c_1 + c_2 t) e^{-\frac{7t}{2}}. \end{cases}$$

$$2. \quad \begin{cases} y = -2c_1 e^{-x} + c_2 e^{-7x}, \\ z = c_1 e^{-x} + c_2 e^{-7x}. \end{cases}$$

$$3. \quad \begin{cases} 2x = e^{-6t}[(c_1 + c_2) \sin t + (c_2 - c_1) \cos t], \\ y = e^{-6t}(c_1 \sin t + c_2 \cos t). \end{cases}$$

$$4. \quad \begin{cases} y = c_1 e^{nx} + c_2 e^{-nx} + \frac{e^x}{n^2 - 1}, \\ az = -nc_1 e^{nx} + nc_2 e^{-nx} - \frac{e^x}{n^2 - 1}. \end{cases}$$

$$5. \quad \begin{cases} x = (c_1 + c_2 t) e^{-4t} - \frac{e^{2t}}{36} + \frac{4e^t}{25}, \\ y = -(c_1 + c_2 + c_2 t) e^{-4t} + \frac{7e^{2t}}{36} + \frac{e^t}{25}. \end{cases}$$

$$6. \quad \begin{cases} x = 2c_1 e^{-4t} - c_2 e^{-7t} + \frac{e^{2t}}{27} + \frac{7e^t}{40}, \\ y = c_1 e^{-4t} + c_2 e^{-7t} + \frac{7e^{2t}}{54} + \frac{e^t}{40}. \end{cases}$$

$$7. \begin{cases} x = c_1 e^{-t} + c_2 e^{-6t} - \frac{29 e^t}{7} + \frac{19 t}{3} - \frac{56}{9}, \\ y = -c_1 e^{-t} + 4 c_2 e^{-6t} + \frac{24 e^t}{7} - \frac{17 t}{3} + \frac{55}{9}. \end{cases}$$

$$8. \begin{cases} x = (c_1 + c_2 t) e^{-4t} - \frac{49 e^{2t}}{36} + \frac{31 e^t}{25}, \\ y = -(c_1 + c_2 + c_2 t) e^{-4t} + \frac{19 e^{2t}}{36} - \frac{11 e^t}{25}. \end{cases}$$

$$9. \begin{cases} x = (c_1 \sin t + c_2 \cos t) e^{-4t} + \frac{31 e^t}{26} - \frac{93}{17}, \\ y = [(c_2 - c_1) \sin t - (c_2 + c_1) \cos t] e^{-4t} - \frac{2 e^t}{13} + \frac{6}{17}. \end{cases}$$

$$10. \begin{cases} x = c_1 t^{-4} + 2 c_2 t^{-3} + \frac{3 t}{10} + \frac{t^2}{15}, \\ y = -c_1 t^{-4} - c_2 t^{-3} - \frac{t}{20} + \frac{2 t^2}{15}. \end{cases}$$

$$11. \begin{cases} x = c_1 t^{-2} + \frac{t}{3}, \\ y = c_2 e^t - c_1 t^{-2} - \frac{t}{3}. \end{cases}$$

$$12. \begin{cases} x = a_1 \sin kt + a_2 \cos kt + a_3, \\ y = b_1 \sin kt + b_2 \cos kt + b_3, \\ z = c_1 \sin kt + c_2 \cos kt + c_3, \end{cases}$$

$$\text{where } k^2 = l^2 + m^2 + n^2.$$

The arbitrary constants are connected by the following equations :

$$\frac{mc_1 - nb_1}{a_2} = \frac{na_1 - lc_1}{b_2} = \frac{lb_1 - ma_1}{c_2} = k,$$

$$la_1 + mb_1 + nc_1 = 0, \quad \frac{\alpha_3}{l} = \frac{b_3}{m} = \frac{c_3}{n}$$

$$13. \begin{cases} x = a_1 \sin (k \log t) + a_2 \cos (k \log t) + a_3, \\ y = b_1 \sin (k \log t) + b_2 \cos (k \log t) + b_3, \\ z = c_1 \sin (k \log t) + c_2 \cos (k \log t) + c_3, \end{cases}$$

where $k^2 = l^2 + m^2 + n^2$.

The arbitrary constants are connected by the following equations :

$$\frac{mn(c_1 - b_1)}{la_2} = \frac{nl(a_1 - c_1)}{mb_3} = \frac{lm(b_1 - a_1)}{nc_2} = k,$$

$$l^2 a_1 + m^2 b_1 + n^2 c_1 = 0, \quad a_2 = b_3 = c_3.$$

Art. 62.

1. $\begin{cases} x = c_1 \sin nt + c_2 \cos nt, \\ y = c_3 + c_4 t - x. \end{cases}$
2. $\begin{cases} x = (c_1 + c_2 t) e^t + (c_3 + c_4 t) e^{-t} - 23, \\ -2y = (c_1 - c_2 + c_2 t) e^t + (c_3 + c_4 + c_4 t) e^{-t} - 36. \end{cases}$
3. $\begin{cases} x = 4c_1 e^{2t} + 4c_2 e^{-2t} + c_3 e^{t\sqrt{7}} + c_4 e^{-t\sqrt{7}} + \frac{1}{7}, \\ y = c_1 e^{2t} + c_2 e^{-2t} + c_3 e^{t\sqrt{7}} + c_4 e^{-t\sqrt{7}} + \frac{9}{14}. \end{cases}$
4. $\begin{cases} x = e^{\frac{nt}{\sqrt{2}}} \left(c_1 \sin \frac{nt}{\sqrt{2}} + c_2 \cos \frac{nt}{\sqrt{2}} \right) + e^{-\frac{nt}{\sqrt{2}}} \left(c_3 \sin \frac{nt}{\sqrt{2}} + c_4 \cos \frac{nt}{\sqrt{2}} \right), \\ y = e^{\frac{nt}{\sqrt{2}}} \left(c_2 \sin \frac{nt}{\sqrt{2}} - c_1 \cos \frac{nt}{\sqrt{2}} \right) + e^{-\frac{nt}{\sqrt{2}}} \left(-c_4 \sin \frac{nt}{\sqrt{2}} + c_3 \cos \frac{nt}{\sqrt{2}} \right). \end{cases}$
5. $\begin{cases} y = (c_1 + c_2 x) e^x + 3c_3 e^{-\frac{3x}{2}} - \frac{x}{2}, \\ z = 2(3c_3 - c_1 - c_2 x) e^x - c_3 e^{-\frac{3x}{2}} - \frac{1}{3}. \end{cases}$
6. $y = u + v, \quad z = -u + v,$
 where $u = c_1 e^{2x} + (c_2 e^{nx} + c_3 e^{-nx}) e^x,$
 $v = c_4 e^{-2x} + (c_5 e^{nx} + c_6 e^{-nx}) e^{-x}.$

7. $y = u + v, \quad z = u - v,$

where $u = (c_1 + c_2x + c_3e^{x\sqrt{2}} + c_4e^{-x\sqrt{2}})e^x,$
 $v = (c_5 + c_6x + c_7e^{x\sqrt{2}} + c_8e^{-x\sqrt{2}})e^{-x}.$

Art. 63.

1. $x = cy^a.$

2. $y^2 = 4ax + c,$ a parabola.

3. $\pm(x + c) = \log(y + \sqrt{y^2 - 1}),$
 or $y = \frac{1}{2}(e^{x+c} + e^{-x-c}),$ a catenary.

4. $4my + c = 2mx\sqrt{4m^2x^2 - 1} + \log(2mx - \sqrt{4m^2x^2 - 1}).$

5. $\pm(x + c) = a \log(y + \sqrt{y^2 - a^2}),$
 or $\pm(x + c) = a \log \frac{y + \sqrt{y^2 - a^2}}{a},$
 from which $y = \frac{a}{2} \left(e^{\frac{x+c}{a}} + e^{-\frac{x+c}{a}} \right),$ a catenary.

6. $x^2 + y^2 = c^2,$ a circle.

7. $x^2 + y^2 - 2cy = 0,$ a circle.

8. $2x^2 + y^2 = 2c^2,$ an ellipse.

9. $x^2 - y^2 = c^2,$ an equilateral hyperbola.

10. $y^2 + x^2 = a^2 \log x^2 + c.$

11. $y = cx^m.$

12. $\frac{x^2}{h^2 - c^2} - \frac{y^2}{c^2} = 1,$ an ellipse or hyperbola.

13. $x^{\frac{1}{2}} + y^{\frac{1}{2}} = a^{\frac{1}{2}},$ a parabola.

14. $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}},$ a hypocycloid.

15. $2xy = a^2,$ an equilateral hyperbola.

16. $c^2y^2 - \log y^2 = 4c(x + c').$

17. $cy^2 - \frac{c^2}{n}(x+c')^2 = 1$, a hyperbola, when $n > 0$; an ellipse or hyperbola, when $n < 0$.

18. First, $(x+c')^2 + y^2 = c^2$, a circle.

Second, $\pm(x+c') = c \log(y + \sqrt{y^2 - c^2})$,

or $\pm(x+c') = c \log \frac{y + \sqrt{y^2 - c^2}}{c}$,

from which $y = \frac{c}{2} \left(e^{\frac{x+c'}{c}} + e^{-\frac{x+c'}{c}} \right)$, a catenary.

19. First, $x+c' = c \operatorname{vers}^{-1} \frac{y}{c} - \sqrt{2cy - y^2}$, a cycloid

Second, $(x+c')^2 = 2cy - c^2$, a parabola.

The Method of Least Squares.

With Numerical Examples of its Application. By GEORGE G. COMSTOCK, Professor of Astronomy in the University of Wisconsin, and Director of the Washburn Observatory. 8vo. Cloth. viii + 68 pages. Mailing price, \$1.05; for introduction, \$1.00.

College Requirements in Algebra: A Final Review.

By GEORGE PARSONS TIBBETS, A.M., Instructor in Mathematics, Williston Seminary. 12mo. Cloth. 46 pages. By mail, 55 cents; to teachers and for introduction, 50 cents.

FROM a wide collection of college papers about 400 examples, illustrating nearly every principle in Algebra, were selected and carefully arranged by subjects.

Peirce's Elements of Logarithms.

With an explanation of the author's Three and Four Place Tables. By Professor JAMES MILLS PEIRCE, of Harvard University. 12mo. Cloth. 80 pages. Mailing price, 55 cents; for introduction, 50 cents.

Mathematical Tables Chiefly to Four Figures.

With full explanations. By Professor JAMES MILLS PEIRCE, of Harvard University. 12mo. Cloth. Mailing price, 45 cents; for introduction, 40 cents.

Elements of the Differential Calculus.

With numerous Examples and Applications. Designed for Use as a College Text-Book. By W. E. BYERLY, Professor of Mathematics, Harvard University. 8vo. 273 pages. Mailing price, \$2.15; for introduction, \$2.00.

THE peculiarities of this treatise are the rigorous use of the Doctrine of Limits, as a foundation of the subject, and as preliminary to the adoption of the more direct and practically convenient infinitesimal notation and nomenclature; the early introduction of a few simple formulas and methods for integrating; a rather elaborate treatment of the use of infinitesimals in pure geometry; and the attempt to excite and keep up the interest of the student by bringing in throughout the whole book, and not merely at the end, numerous applications to practical problems in geometry and mechanics.

Elements of the Integral Calculus.

Second Edition, revised and enlarged. By W. E. BYERLY, Professor of Mathematics in Harvard University. 8vo. xvi + 383 pages. Mailing price, \$2.15; for introduction, \$2.00.

THIS work contains, in addition to the subjects usually treated in a text-book on the Integral Calculus, an introduction to Elliptic Integrals and Elliptic Functions; the Elements of the Theory of Functions; a Key to the Solution of Differential Equations; and a Table of Integrals.

John E. Clark, *Prof. of Mathematics, Sheffield Scientific School of Yale University*: The additions to the present edition seem to me most judicious and to greatly enhance its

value for the purposes of university instruction, for which in several important respects it seems to me better adapted than any other American text-book on the subject.

An Elementary Treatise on Fourier's Series,

and Spherical, Cylindrical, and Ellipsoidal Harmonics, with Applications to Problems in Mathematical Physics.

By WILLIAM E. BYERLY, Ph.D., Professor of Mathematics, Harvard University. 8vo. Cloth. x + 288 pages. Mailing price, \$3.15; for introduction, \$3.00.

THIS book is intended as an introduction to the treatment of some of the important Linear Partial Differential Equations which lie at the foundation of modern theories in physics, and deals mainly with the methods of building up solutions of a differential equation from easily obtained particular solutions, in such a manner as to satisfy given initial conditions.

John Perry, *Technical College, Finsbury, London, England*: Byerly's book is one of the most useful books in existence. I have read it with great delight and I am happy

to say that although it seemed to be written expressly for me, one of my friends who is a great mathematician, seems as delighted with it as I am myself.

A Short Table of Integrals. *Revised and Enlarged.*

By B. O. PEIRCE, Prof. Math., Harvard Univ. 32 pages. Mailing price, 15 cents. Bound also with Byerly's Calculus.

Byerly's Syllabi.

Each, 8 or 12 pages, 10 cents. The series includes,—Plane Trigonometry, Plane Analytical Geometry, Plane Analytic Geometry (*Advanced Course*), Analytical Geometry of Three Dimensions, Modern Methods in Analytic Geometry, the Theory of Equations.

Academic Trigonometry: Plane and Spherical.

By T. M. BLAKSLÉE, Ph.D. (Yale), Professor of Mathematics in Des Moines College, Iowa. 12mo. Cloth. 33 pages. Mailing price, 30 cents; for introduction, 25 cents.

THE Plane and Spherical portions are arranged on opposite pages.

The memory is aided by analogies, and it is believed that the entire subject can be mastered in less time than is usually given to Plane Trigonometry alone, as the work contains but 29 pages of text.

The Plane portion is compact, and complete in itself.

Examples of Differential Equations.

By GEORGE A. OSBORNE, Professor of Mathematics in the Massachusetts Institute of Technology, Boston. 12mo. Cloth. vii + 50 pages. Mailing Price, 60 cents; for introduction, 50 cents.

A SERIES of nearly three hundred examples with answers, systematically arranged and grouped under the different cases, and accompanied by concise rules for the solution of each case.

Selden J. Coffin, *Prof. of Astronomy*, *Lafayette College*: Its appearance is most timely, and it supplies a manifest want.

Determinants.

The Theory of Determinants: an Elementary Treatise. By PAUL H. HANUS, B.S., recently Professor of Mathematics in the University of Colorado, now Assistant Professor, Harvard University. 8vo. Cloth. viii + 217 pages. Mailing price, \$1.90; for introduction, \$1.80.

THIS book is written especially for those who have had no previous knowledge of the subject, and is therefore adapted to self-instruction as well as to the needs of the class-room. The subject is at first presented in a very simple manner. As the reader advances, less and less attention is given to details. Throughout the entire work it is the constant aim to arouse and enliven the reader's interest, by first showing how the various concepts have arisen naturally, and by giving such applications as can be presented without exceeding the limits of the treatise.

William G. Peck, *late Prof. of Mathematics, Columbia College, N. Y.*: A hasty glance convinces me that it is an improvement on Muir.

T. W. Wright, *Prof. of Mathematics, Union Univ., Schenectady, N. Y.*: It fills admirably a vacancy in our mathematical literature, and is a very welcome addition indeed.

Analytic Geometry.

By A. S. HARDY, Ph.D., Recently Professor of Mathematics in Dartmouth College, and author of *Elements of Quaternions*. 8vo. Cloth. xiv + 239 pages. Mailing price, \$1.60; for introduction, \$1.50.

THIS work is designed for the student, not for the teacher. Particular attention has been given to those fundamental conceptions and processes which, in the author's experience, have been found to be sources of difficulty to the student in acquiring a grasp of the subject as a *method of research*. The limits of the work are fixed by the time usually devoted to Analytic Geometry in our college courses by those who are not to make a special study in mathematics. It is hoped that it will prove to be a *text-book* which the teacher will wish to use in his class-room, rather than a *book of reference* to be placed on his study shelf.

Oren Root, Professor of Mathematics, Hamilton College: It meets quite fully my notion of a text for our classes. I have hesitated somewhat about introducing a generalized discussion of the conic in required work. I have, however, read Mr. Hardy's discussion carefully twice; and it seems to me that a student who can get the subject at all can get that. It is my present purpose to use the work next year.

John E. Clark, Professor of Mathematics, Sheffield Scientific School of Yale College: I need not hesitate to say, after even a cursory examination, that it seems to me a very attractive book, as I anticipated it

would be. It has evidently been prepared with real insight alike into the nature of the subject and the difficulties of beginners, and a very thoughtful regard to both; and I think its aims and characteristic features will meet with high approval. While leading the student to the usual useful results, the author happily takes especial pains to acquaint him with the character and spirit of analytical methods, and, so far as practicable, to help him acquire skill in using them.

John B. French, Dean of College of Liberal Arts, Syracuse University: It is a very excellent work, and well adapted to use in the recitation room.

Elements of Quaternions.

By A. S. HARDY, Ph.D., Recently Professor of Mathematics in Dartmouth College. *Second edition, revised*. Crown. 8vo. Cloth. viii + 234 pages. Mailing price, \$2.15; for introduction, \$2.00.

THE chief aim has been to meet the wants of beginners in the class-room, and it is believed that this work will be found superior in fitness for beginners in practical compass, in explanations and applications, and in adaptation to the methods of instruction common in this country.

Elements of the Calculus.

By A. S. HARDY, Ph.D., Recently Professor of Mathematics in Dartmouth College. 8vo. Cloth. xi + 239 pages. Mailing price, \$1.60; for introduction, \$1.50.

THIS text-book is based upon the method of rates. The object of the Differential Calculus is the measurement and comparison of rates of change when the change is not uniform. Whether a quantity is or is not changing uniformly, however, its rate at any instant is determined essentially in the same manner, viz. : by letting it change at the rate it had at the instant in question and observing what this change is. It is this change which the Calculus enables us to determine, however complicated the law of variation may be. From the author's experience in presenting the Calculus to beginners, the method of rates gives the student a more intelligent, that is, a less mechanical, grasp of the problems within its scope than any other. No comparison has been made between this method and those of limits and of infinitesimals. This larger view of the Calculus is for special or advanced students, for which this work is not intended; the space and time which would be required by such general comparison being devoted to the applications of the method adopted.

Part I., Differential Calculus, occupies 166 pages. Part II., Integral Calculus, 73 pages.

George B. Merriman, *formerly Prof. of Mathem. and Astron. Rutgers College*: I am glad to observe that Professor Hardy has adopted the method of rates in his new Calculus, a logical and intelligent method, which avoids certain difficulties involved in the usual methods.

J. B. Coit, *Prof. of Mathematics, Boston University*: It pleases me very much. The treatment of the first principles of Calculus by the method of rates is eminently clear. Its use next year is quite probable.

Ellen Hayes, *Prof. of Mathematics, Wellesley College*: I have found

it a pleasure to examine the book. It must commend itself in many respects to teachers of Calculus.

W. R. McDaniel, *Prof. of Mathematics, Western Maryland College*: Hardy's Calculus and Analytic Geometry are certainly far better books for the college class-room than any others I know of. The feature of both books is the directness with which the author gets right at the very fact that he intends to convey to the student, and the force of his presentation of the fact is greatly augmented by the excellent arrangement of type and other features of the mechanical make-up.